

Lesson 7-1

Example 1 Find Degree and Leading Coefficients

State the degree and leading coefficient of each polynomial in one variable. If it is not a polynomial in one variable, explain why.

a. $-2a^3 + a - \frac{1}{a^2}$

This is not a polynomial. The term $\frac{1}{a^2}$ cannot be written in the form x^n , where n is a nonnegative integer.

b. $b^5 + b^2 - 8$

This is a polynomial in one variable. The degree is 5, and the leading coefficient is 1.

c. $mn + 3m - 7n$

This is not a polynomial in one variable. It contains two variables, m and n .

d. $11 - 5c + 2c^6 - 4c^3$

Rewrite the expression so the powers of c are in decreasing order.

$$2c^6 - 4c^3 - 5c + 11$$

This is a polynomial in one variable. The degree is 6, and the leading coefficient is 2.





Example 2 Evaluate a Polynomial Function

GEOMETRY In a polygon with n sides, for $n > 3$, the number of diagonals can be found by using

the function $d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$, where $d(n)$ is the number of diagonals and n is the number of sides.

a. Draw polygons for $n = 4, 5, 6$, and 7 . Show that the polynomial function $d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$ gives the total number of diagonals for each figure.

First, draw the polygons and count the number of diagonals.



Next, find the values of $d(4)$, $d(5)$, $d(6)$, and $d(7)$.

$$d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$$

$$\begin{aligned}d(4) &= \frac{1}{2}(4)^2 - \frac{3}{2}(4) \\ &= 8 - 6 \\ &= 2\end{aligned}$$

$$d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$$

$$\begin{aligned}d(5) &= \frac{1}{2}(5)^2 - \frac{3}{2}(5) \\ &= 12.5 - 7.5 \\ &= 5\end{aligned}$$

$$d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$$

$$\begin{aligned}d(6) &= \frac{1}{2}(6)^2 - \frac{3}{2}(6) \\ &= 18 - 9 \\ &= 9\end{aligned}$$

$$d(n) = \frac{1}{2}n^2 - \frac{3}{2}n$$

$$\begin{aligned}d(7) &= \frac{1}{2}(7)^2 - \frac{3}{2}(7) \\ &= 24.5 - 10.5 \\ &= 14\end{aligned}$$

You can see that the number of diagonals for each figure in the diagrams is the same as the number of diagonals $d(n)$ found by evaluating the function.

b. Find the number of diagonals for a decagon (a ten-sided polygon).

$$\begin{array}{ll}
 d(n) = \frac{1}{2}n^2 - \frac{3}{2}n & \text{Given function} \\
 d(10) = \frac{1}{2}(10)^2 - \frac{3}{2}(10) & \text{Replace } n \text{ with } 10. \\
 = 50 - 15 & \text{Evaluate.} \\
 = 35 & \text{Simplify.}
 \end{array}$$

Example 3 Functional Values of Variables

a. Find $p(-2)$ if $p(x) = -x^4 - x^3 + 6$.

$$\begin{array}{ll}
 p(x) = -x^4 - x^3 + 6 & \text{Original function} \\
 p(-2) = -(-2)^4 - (-2)^3 + 6 & \text{Replace } x \text{ with } -2. \\
 = -16 - (-8) + 6 & \text{Simplify.} \\
 = -2 & \text{Simplify.}
 \end{array}$$

b. Find $-3[q(4b - 1)]$ if $q(x) = x^2 - x + 1$.

First, evaluate $q(4b - 1)$ by replacing x in $q(x)$ with $4b - 1$.

$$\begin{array}{ll}
 q(x) = x^2 - x + 1 & \text{Original function} \\
 q(4b - 1) = (4b - 1)^2 - (4b - 1) + 1 & \text{Replace } x \text{ with } 4b - 1. \\
 = 16b^2 - 8b + 1 - 4b + 1 + 1 & \text{Evaluate } (4b - 1)^2 \text{ and } -1(4b - 1). \\
 = 16b^2 - 12b + 3 & \text{Simplify.}
 \end{array}$$

To evaluate $-3[q(4b - 1)]$, multiply $q(4b - 1)$ by -3 .


$$\begin{array}{ll}
 -3[q(4b - 1)] = -3(16b^2 - 12b + 3) & \text{Substitute } 16b^2 - 12b + 3 \text{ for } q(4b - 1). \\
 = -48b^2 + 36b - 9 & \text{Distributive Property}
 \end{array}$$

Example 4 Graphs of Polynomial Functions


For each graph:

- Describe the end behavior.
- Determine whether it represents an odd-degree or an even-degree polynomial function.
- State the number of real zeros.


a.



b.



c.



- a.
- As $x \rightarrow +\infty, f(x) \rightarrow +\infty$. As $x \rightarrow -\infty, f(x) \rightarrow +\infty$.
 - It is an even-degree polynomial function.
 - The graph does not intersect the x -axis, so it has no real zeros.
- b.
- As $x \rightarrow +\infty, f(x) \rightarrow +\infty$. As $x \rightarrow -\infty, f(x) \rightarrow -\infty$.
 - It is an odd-degree polynomial function.
 - The graph intersects the x -axis at three points, so the function has three real zeros.
- c.
- As $x \rightarrow -\infty, f(x) \rightarrow +\infty$. As $x \rightarrow +\infty, f(x) \rightarrow -\infty$.
 - It is an odd-degree polynomial function.
 - The graph intersects the x -axis at one point, so the function has one real zero.